

12:47



[Test results](#)

**Test detail**



Demographics

Age: 65 y.o.

Gender: female



Medical/Surgical History

Menstrual periods:None

Age of menopause:57

History of hip fractures in parents:No

History of previous fractures occurring spontaneously, or a fracture as a result of a fall from a standing height or less:No

If yes, indicated area:

Glucocorticoids use:No

Currently taking medication:None

Have or had listed medical conditions:Hyperparathyroidism

Dietary/Habit

Alcohol 3units or more per day (on average):No

Currently smoking:No

Other pertinent history:COMPARISON

Dual-X-ray Absorptiometry (DXA Results)

AP spine (L1-L4) Left hip (neck) Left hip (total) Left forearm (1/3)

BMD (gm/cm<sup>2</sup>) 0.846 0.538 0.621 0.598

T score -1.8 -2.8 -2.6 -1.6

Z score 0.0 -1.3 -1.4 0.1

%change from previous scan dated: 12-07-2015 -1.7% -3.7%

FRAX Score:

10 YR Risk Hip Fracture % na

Typical threshold to start therapy in patients is a 10-year risk of hip fracture >3%. Clinician's judgment and/or patient preferences may indicate treatment for people with 10-year fracture probabilities above or below these levels.

10 YR Risk Major Osteoporotic Fracture % na

Typical threshold to start therapy in patients is a 10-year risk of any osteoporotic fracture >20%. Clinicians judgment and/or patient preferences may indicate treatment for people with 10-year fracture probabilities above or below these levels.

Diagnosis:



Home



My Care



My Plan



Find Care



Inbox

138